

Cultured Adherent Cell Sample Preparation for the RNAscope® Multiplex Fluorescent v2 Assay

Introduction

This Technical Note provides guidelines for the preparation of cultured adherent cells that can be assayed using the RNAscope® Multiplex Fluorescent Kit v2 (Cat. No. 323100). To pretreat the cells, use RNAscope® Protease III (available in the RNAscope® Protease III and Protease IV Reagents, Cat. No. 322340 or RNAscope® Universal Pretreatment Kit,

Cat No 322380). Read the Safety Data Sheet (SDS) available on the website and follow handling instructions. Wear appropriate protective eyewear, clothing, and gloves. For the latest services and support information, go to: www.acdbio.com/support.

Workflow

Part 1: Cell Collection

Cell Culture

1. One day before fixation, seed cells in growth medium on chamber slides at a density that will allow cells to be 80–90% confluent at the time of fixation.

Cell Fixation

1. Remove growth media, and disassemble chambers.
2. Submerge the slides in a Tissue-Tek® container with 200 mL of 1X PBS.

IMPORTANT! Do not let cells dry out at any time. Always use enough solution to submerge all the cells.

3. Remove 1X PBS, and add 10% Neutral Buffered Formalin (NBF). Incubate at **ROOM TEMPERATURE (RT)** for **30 MIN**.
4. Remove NBF, and gently rinse slides with 1X PBS. Repeat twice.

Dehydrate and Store Cells

1. Remove 1X PBS wash, and replace with 50 mL 50% EtOH. Incubate at **RT** for **5 MIN**.
2. Remove 50% EtOH, and replace with 50 mL 70% EtOH. Incubate at **RT** for **5 MIN**.
3. Remove 70% EtOH, and replace with 50 mL 100% EtOH. Incubate at **RT** for **5 MIN**.
4. Remove 100% EtOH, and replace with fresh 100% EtOH. Incubate at **RT** for **10 MIN**.

NOTE: The slides can be stored in 100% EtOH at **-20°C** for up to **6 MONTHS**.

Part 2: Cell Pretreatment

Rehydrate Cells

1. Submerge slides in 70% EtOH. Incubate at **RT** for **2 MIN**.

IMPORTANT! Do not let cells dry out at any time. Always use enough solution to submerge all the cells.

2. Remove 70% EtOH, and replace with 50% EtOH. Incubate at **RT** for **2 MIN**.
3. Remove 50% EtOH, and replace with 1X PBS. Incubate at **RT** for **10 MIN**.

Apply RNAscope® Hydrogen Peroxide

1. Lay slides on the bench, and add ~5–8 drops of RNAscope® Hydrogen Peroxide to cover the entire section.
2. Incubate the slides at **RT** for **10 MIN**.
3. Remove the RNAscope® Hydrogen Peroxide solution from one slide at a time by tapping and/or flicking the slide on absorbent paper. Immediately insert the slide into a Tissue-Tek® Slide Rack submerged in a Tissue-Tek® Staining Dish filled with distilled water.
4. Wash slides 3–5 times by moving the Tissue-Tek® Slide Rack up and down in the distilled water.
5. Repeat Step 4 with fresh distilled water.

Create a Hydrophobic Barrier

1. Draw 2–4 times around each well/circle on the chambered slides using the Immedge™ hydrophobic barrier pen. Let the barrier dry completely **~1 MIN**.

NOTE: Do not let the cells dry out during this step. Place slides back into 1X PBS if the cells look too dry.

2. Rinse slides briefly with 1X PBS in a Coplin jar or staining dish.

Apply RNAscope® Protease III

1. One at a time, remove each slide from the 1X PBS and tap/flick to remove excess liquid. Place the

slides on the HybEZ™ Slide Rack, and place rack in the Humidity Control Tray.

2. Add 2–4 drops diluted Protease III to completely cover each well/circle.

NOTE: For most cell lines, dilute protease **1:15** with 1X PBS. Protease dilution factor must be empirically determined for each new cell type.

3. Close the Humidity Control Tray and incubate for **10 MIN** at **RT**.
4. One at a time, take each slide from the HybEZ™ Slide Rack and tap/flick to remove excess liquid. Submerge slides in 1X PBS.
5. Wash the slides by agitating them in the 1X PBS. Repeat with fresh 1X PBS.

IMPORTANT! Proceed to the RNAscope® protocol using the *RNAscope® Multiplex Fluorescent Kit User Manual Part 2* (Catalog No. 323100) available at <https://acdbio.com/technical-support/user-manuals>.

Obtaining Support

For the latest services and support information, go to: <https://acdbio.com/technical-support>.

At the website, you can:

- Access telephone and fax numbers to contact Technical Support and Sales.
- Search through FAQs.
- Submit a question directly to Technical Support.

For Research Use Only. Not for diagnostic use.

NOTICE TO PURCHASER: PLEASE REFER TO THE RNASCOPE® FLUORESCENT MULTIPLEX V2 ASSAY- USER MANUAL FOR LIMITED USE LABEL LICENSE OR DISCLAIMER INFORMATION. Advanced Cell Diagnostics, Inc. reserves the right to change its products and services at any time to incorporate technological developments. This manual is subject to change without notice. Although this manual has been prepared with every precaution to ensure accuracy, Advanced Cell Diagnostics, Inc. assumes no liability for any errors, omissions, or for any damages resulting from the use of this information.

© 2017 Advanced Cell Diagnostics. All rights reserved. RNAscope® and HybEZ™ are trademarks of Advanced Cell Diagnostics, Inc. All other trademarks belong to their respective owners.

Headquarters

7707 Gateway Blvd Suite 200, Newark, CA 94545 Phone 1-510-576-8800 Toll Free 1-877-576-3636

For support, email support@acdbio.com

www.acdbio.com

